arXiv:1209.3128v1 [math.AG] 14 Sep 2012

METRICS ON UNIVERSAL COVERING OF PROJECTIVE VARIETY

Robert Treger

1. Preliminaries

Unless stated otherwise, throughout the note $X \hookrightarrow \mathbf{P}^r$ will be an nonsingular connected projective variety of dimension n with *large* and *residually finite* fundamental group $\pi_1(X)$. Let U_X denote its universal covering.

Let $\mathcal{L}_X$ be a very ample line bundle on X defining the embedding. Given X and $\mathcal{L}_X$ as above, we will define a real analytic $\pi_1(X)$-invariant Kahler metric Λ on U_X. The metric will be given by Calabi's diastatic potential [C].

Next, we establish another version of the uniformization theorem [T1] by constructing yet another $\pi_1(X)$-invariant Kahler metric Σ on U_X provided $\pi_1(X)$ is nonamenable. It will be a Bergman-type metric with negative Ricci form. Hence the canonical bundle $\mathcal{K}_X$ will be ample by Kodaira. So, we can apply our earlier uniformization [T1, Sections 3, 4] because $\pi_1(X)$ is nonamenable.

Diastasis [C]**.** Let M denote a complex manifold with real analytic Kahler metric. Let Φ denote a real analytic potential of the metric defined in a small neighborhood $\mathcal{V} \subset M$. Then there exists a unique holomorphic functional element F on an open neighborhood of the diagonal in $M \times \bar{M}$ such that $F(p,p) = \Phi(p)$. Then the functional element of diastasis is defined by

$$D_M(p,q) := F(p,p) + F(q,q) - F(p,q) - F(q,p) \quad (p, q \in \mathcal{V}).$$

2. Metric Λ

The hyperplane bundle on $\mathbf{P}^r$ restricts to $\mathcal{L}_X$, called a polarization on X. Given any polarized Kahler metric g on X, one can find a Hermitian metric h on $\mathcal{L}_X$. We will consider the volume form dV_g, and we will use the same volume form on all the coverings of X. For each positive integer m_0, the Hermitian metric h induces a Hermitian metric h^{m_0} on $\mathcal{L}_X^{m_0}$, as well as on all inverse images of $\mathcal{L}_X^{m_0}$ on the coverings of X.

One chooses an orthonormal basis $(s_0^{m_0}, \dots, s_{r_{m_0}}^{m_0})$ of the space $H^0(X, \mathcal{L}_X^{m_0})$ of all global sections of $\mathcal{L}_X^{m_0}$. We get an inner product and a natural embedding:

$$\langle s_\alpha^{m_0}, s_\beta^{m_0} \rangle := \frac{1}{Vol_g(X)} \int_X h^{m_0}(s_\alpha^{m_0}, s_\beta^{m_0}) dV_g; \qquad \phi_{X,m_0} : X \hookrightarrow \mathbf{P}^{r_{m_0}}.$$

Typeset by $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-TEX

Let g_{FS} be the standard Fubini-Study metric on $P^{r_{m_0}}$. The $\frac{1}{m_0}$-multiple of g_{FS} on $P^{r_{m_0}}$ restricts to a Kahler metric on X:

$$g_{X,m_0} := \frac{1}{m_0}\phi^*_{X,m_0} g_{FS}.$$

One of the main results of Tian in [Ti] (Yau's conjecture) says g_{X,m_0} converges to g_X in C^2-topology. His proof is local by nature.

Similar statement holds for all finite coverings of X as well as for a small neighborhood $\mathcal{V}_p$ of a point $p \in \mathcal{V}_p \subset U_X$. One considers L^2-integrable sections over $\mathcal{V}_p$. We consider a tower of Galois coverings with each $Gal(X_i/X)$ a finite group:

$$X = X_0 \leftarrow X_1 \leftarrow X_2 \leftarrow \cdots \leftarrow U_X, \quad \bigcap_i Gal(U_X/X_i) = \{1\} \quad (0 \le i < \infty).$$

Let $\tau_i : X_i \to X$ and $\tau : U_X \to X$ denote the projections. Then the bundles $\tau_i^*\mathcal{L}_X$ $(0 \le i < \infty)$ are ample. However, $\tau_i^*\mathcal{L}_X$'s are not necessary very ample bundles.

For an appropriate m_i, the bundle $(\tau_i^*\mathcal{L}_X)^{\otimes m_i}$ is very ample hence it defines an embedding $\phi_{X_i,m_i} : X_i \hookrightarrow \mathbf{P}^{r_{m_i}}$. As above, we get a metric g_{X_i,m_i} on X_i and the corresponding diastatic potential.

Finally, we consider the integers $m_{ij} := m_i + j$ for $0 \le i, j < \infty$. As above, we obtain the metrics $g_{X_i,m_{ij}}$ on X_i.

As we mentioned, the volume forms are arising from X, and the Hermitian metrics on the corresponding bundles are arising from the Hermitian metric h on $\mathcal{L}_X$. The pullbacks to $\mathcal{V}_p$ of the metrics on X_i's are *smaller* than the corresponding metrics on $\mathcal{V}_p$.

We pull back all the metrics in the tower of *finite* coverings of X to the neighborhood $\mathcal{V}_p$. By the result of Tian [Ti], we get the converging sequence of metrics in the neighborhood $\mathcal{V}_p$. All the metrics in the sequence are the Bergman-type metrics. Hence, we get the convergence of the corresponding *Hermitian kernels of positive type* (see the definition in [FK, p. 6]). We get the convergence of the diastatic potentials. As the limit, we obtain the diastatic potential of the desired Kahler metric Λ at the point p. It is easy to see that the diastasises generate a global strictly plurisubharmonic *function* on U_X (compare [T2]).

Example. Let C be a curve of genus $g(C) \ge 2$. We will assume:

$$\mathcal{K}_C^{\otimes \ell} \subset \mathcal{L}_C \subset \mathcal{K}_C^{\otimes m}$$

where $\mathcal{K}_C$ is the canonical bundle, and ℓ, m are appropriate large integers. We have the embeddings of C and $\Delta = U_C$ in the corresponding projective spaces given by the corresponding powers of canonical bundles. We get the Bergman metrics on C and Δ. The corresponding Bergman kernels $\mathbf{B}_{\Delta,\mathcal{K}^{\otimes t}}$ are well known for all $t \ge 1$. The corresponding diastatic potential equals $\log \mathbf{B}_{\Delta,\mathcal{K}^{\otimes t}}$.

Now, if C is a general curvilinear section of the projective variety X then we consider the inverse image of C on U_X. We obtain an open Riemann surface $R_C \subset U_X$ in place of the disk Δ. The corresponding Bergman kernel $\mathbf{B}_{R_C,\mathcal{K}^{\otimes t}}$ is the Poincaré series of $\mathbf{B}_{\Delta,\mathcal{K}^{\otimes t}}$ (see, for example, [D, Section 1]).

3. UNIFORMIZATION

In 2010, Frédéric Campana asked the author whether one can establish the uniformization theorem (see [T1]) with assumptions on the fundamental group only. In fact, one can prove the following

Theorem. *Let $X \hookrightarrow \mathbf{P}^r$ will be an arbitrary nonsingular connected projective variety of dimension n with large and residually finite fundamental group $\pi_1(X)$. If $\pi_1(X)$ is nonamenable then U_X is a bounded domain in $\mathbf{C}^n$.*

Proof. Since $\pi_1(X)$ is nonamenable, the space of bounded harmonic functions on U_X is infinite-dimensonal (see [Ki, Theorem C′], [To, Theorem]); moreover, if $\pi_1(X)$ is amenable then the above space contains only constants. Their theorems generalize an earlier result in [LS, Theorems 3, 3′].

The following theorem is proved in [Ki]: *Let Γ be a discrete group acting on a separable metric space M so that the quotient M/Γ is compact. Suppose that H is a linear subspace of the space of all bounded continuous function on M such that no nonconstant function from H may attain its supremum in a point of M. If H is invariant under the action of all operators T_γ, $\gamma \in \Gamma$, given by $T_\gamma f(x) = f(\gamma x)$ ($f \in H, x \in M$), then either H contains only constants or H is infinite-dimensional.*

We proceed by induction on dimension. If $n = 1$ then our theorem follows from the classical theorem of Kobe and Poincaré since $\pi_1(X)$ is nonamenable provided $g(X) \geq 2$. We will assume $n = 2$ and proceed to the uniformization. For $n \geq 2$, the induction step from n to $n + 1$ is similar only easier.

Let $C \subset U_X$ be the preimage of a sufficiently general curvilinear section of X. In local coordinates $z = (z_1, \dots, z_n)$, C is given by $\{f(z) = 0\}$. A global section of $\tau^*\mathcal{L}_X \otimes \mathcal{K}_{U_X}$ can be written locally as $\omega := \frac{q(z)(dz_1\wedge\cdots\wedge dz_n)}{f(z)}$ with $q(z)$ a harmonic function.

One can define the Poincaré residue map $\tau^*\mathcal{L}_X \otimes \mathcal{K}_{U_X} \to \mathcal{K}_C$ [GH, p. 147]:

$$\omega := \frac{q(z)(dz_1 \wedge \cdots \wedge dz_n)}{f(z)} \longmapsto \omega' := (-1)^{i-1}\frac{q(z)(dz_1 \wedge \cdots \wedge \widehat{dz_i} \wedge \cdots \wedge dz_n)}{\frac{\partial f}{\partial z_i}}\Big|_{f=0},$$

where i is such that $\frac{\partial f}{\partial z_i} \neq 0$. Similarly we get the map $\tau^*\mathcal{L}_X^{\otimes t} \otimes \mathcal{K}_{U_X}^{\otimes t} \to \mathcal{K}_C^{\otimes t}$.

First, we will show that how to get an immersion of U_X into infinite-dimensional projective space with its Fubini-Study metric (compare [Kob, Chapter 4.10], [G, Section 3]). We fix a sufficiently large integer t and consider the Hilbert space of square-integrable harmonic sections

$$\mathcal{H} := \big\{\omega \in H^0(\tau^*\mathcal{L}_X^{\otimes t} \otimes \mathcal{K}_{U_X}^{\otimes t}) \mid \int_{U_X} \frac{|q|^2 dVol_\Lambda}{\rho_\Lambda^2} < \infty\big\}$$

where $dVol_\Lambda := \rho_\Lambda d\mu$ is the volume form of the metric Λ and $d\mu$ is the Lebesque measure on U_X. This Hilbert space is defined correctly (compare [Kol, Chapters 5.13, 7.1]). Furthermore $\int_{U_X} \frac{dVol_\Lambda}{\rho_\Lambda^2} < \infty$.

We get a natural map ν into an infinite-dimensional projective space. We will show that ν is an immersion. For a direction ℓ on U_X at the point a, we have to find a section ω_a such that the corresponding $q_a(a) = 0$ and $\frac{\partial q_a}{\partial \ell}(a) \neq 0$.

We assume ℓ is tangent to a curve R (= connected Riemann surface) that is the preimage of a curvilinear section of X. Clearly we can find a nontrivial *bounded* harmonic function that will vanish at a.

If necessary we differentiate this function an appropriate number of times in the direction ℓ at $a \in U_X$. On R, we obtain a bounded function times $\lambda^s(u) := (1-|u|^2)^{-s}$. This follows, for example, from a local result in [DP]. The resulting section ω_a will be in $\mathcal{H}$ because of the definition of Λ. Indeed, $\lambda^2(u) = \pi B_\Delta(u,u)$ where $B_\Delta(u,u)$ is the Bergman kernel function of the disk. For the corresponding $q_a(u)$, we get $q_a(a) = 0$ and $\frac{\partial q_a}{\partial \ell}(a) \neq 0$.

Note. Let ℓ be a real direction on R at a point $a \in R$. Since harmonic functions are real analytic, one can choose a nontrivial bounded harmonic function q_a on U_X and local real coordinates around a on R such that not only $q_a(a) = 0$ but $\frac{\partial q_a}{\partial \ell}(a) \neq 0$. Here ℓ will coincide with one of the local coordinates.

By the above, we obtain a *Bergman-type* Kahler metric on U_X via the immersion in an infinite-dimensional projective space (compare [Kob, Chapter 4.10], [G, Section 3]). This metric, denoted by Σ, will have a negative Ricci form. Moreover Σ will be $\pi_1(X)$-invariant. Hence the canonical bundle on X will be ample by Kodaira.

Next, for an integer t, we can assume that $\mathcal{K}_X^{\otimes t}$ will be very ample on X, and its preimages on all finite coverings of X (in our tower of coverings) will be very ample as well [Kol, Chapter 16. 5].

On all the finite coverings of X, we consider the Bergman-type metrics similar to the Bergman-type metric defined by the Hilbert space $\mathcal{H}$. We, now, consider only holomorphic sections of $\mathcal{K}_X^{\otimes t}$ and $\tau_i^*\mathcal{K}_X^{\otimes t}$.

We pull back on U_X all the metrics and take the limit. The limit will exist because each pullback metric will be smaller than the metric defined by $\mathcal{H}$ (see [T3, 2.1]). Moreover, the limiting metric will a Bergman-type metric. It will define an embedding into an infinite-dimensional projective space.

We conclude the prove of the uniformization in our case exactly as in [T1, Sections 3, 4].

References

[C] E. Calabi, *Isometric imbedding of complex manifolds*, Ann. of Math. **58** (1953), 1–23.

[D] D. Drasin, *Cusp forms and Poincaré series*, Amer. J. Math. **90** (1968), 356–365.

[DP] S. Dai and Y. Pan, *Note on Schwarz-Pick estimates for bounded and positive real part analytic functions*, Proc. Aner. Math. Soc. **136** (2008), 635-640.

[FK] J. Faraut, S. Kaneyuki, A. Korányi, Q.-k. Lu, G. Roos, *Analysis and geometry on complex homogeneous domains*, Birkhäuser, Boston, 2000.

[G] M. Gromov, *Kahler hyperbolicity and L_2-Hodge theory*, J. Differential Geometry **33** (1991), 263-292.

[GH] P. Griffiths, J. Harris, *Principles of Algebraic Geometry*, John Wily, New York, 1978.

[Ki] Y. Kifer, *Bounded harmonic functions on nonamenable covers of compact manifolds*, Isr. J. Math. **61** (1988), 170–178.

[Kob] S. Kobayashi, *Hyperbolic Complex Spaces*, Springer, 1998.

[Kol] J. Kollár, *Shafarevich maps and automorphic forms*, Princeton Univ. Press, Princeton, 1995.

[LS] T. Lyons and D. Sullivan, *Bounded harmonic functions on coverings*, J. Differential Geometry **19** (1984), 299-323.

[Ti] G. Tian, *On a set of polarized Kahler metrics on algebraic manifolds*, J. Differential Geometry **32** (1990), 99–130.

[To] D. Toledo, *Bounded harmonic functions on coverings*, Proc. Amer. Math. Soc. **104** (1988), 1218-1219.

[T1] R. Treger, *Uniformization*, arXiv:math.AG/1001.1951v4.

[T2] ———, *Remark on a conjecture of Shafarevich*, arXiv:math.AG/1008.1745v2.

[T3] ———, *Bergman type metrics in tower of coverings*, arXiv:math.AG/1107.5713v2.

PRINCETON, NJ 08540
E-mail address: `roberttreger117@gmail.com`